



PROTEIN BIOTECHNOLOGIES

CATALOG NUMBER: T1-6X-2

HUMAN LUNG TISSUE LYSATE COLLECTION / FEMALE / MIXED CANCER SUBTYPES

Extract I.D.	Diagnosis	Grade	Stage	TNM	Sex	Age
T1-045-T-1	Adenocarcinoma	2	II	T ₂ N ₀ M _x	F	51
T1-045-N-1	Matched normal tissue				F	51
T1-050-T-1	Adenocarcinoma	3	n/a	n/a	F	70
T1-050-N-1	Matched normal tissue				F	70
T1-055-T-1	Large Cell Carcinoma	n/a	n/a	T ₃ N ₁ M _x	F	60
T1-055-N-1	Matched normal tissue				F	60
T1-056-T-1	Small Cell Carcinoma	n/a	II	n/a	F	56
T1-056-N-1	Matched normal tissue				F	56
T1-057-T-1	Spindle Cell Carcinoma	n/a	I	T ₂ N ₀ M ₀	F	25
T1-057-N-1	Matched normal tissue				F	25
T1-058-T-1	Papillary Carcinoma	1	IIIA	T ₂ N ₀ M ₀	F	41
T1-058-N-1	Matched normal tissue				F	41

Concentration: 1 mg/ml, 100 µg/vial.

The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides.

Storage: Aliquot single use volumes to avoid repeated freeze/thaw cycles.

From time of receipt, this product is stable for 3 months at -20°C, or 12 months at -70°C.

Lysate Preparation:

Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Modified RIPA Buffer:

PBS, pH 7.4	1 µg/ml Aprotinin	1 mM NaF
1 mM EDTA	1 µg/ml Pepstatin-A	0.1% SDS
0.25% Na deoxycholate	1 µg/ml Leupeptin	1 mM PMSF
1 mM Na ₃ VO ₄		

Application:

These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required.

Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays.

This material has tested negative for HbsAg, HIV 1/2, and HCV. Use *UNIVERSAL PRECAUTIONS* when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately.

Source:

Integrated Laboratory Services-Biotech (ILSbio), Chestertown, MD 21620 www.ilsbio.com

For Research Use Only