



PROTEIN BIOTECHNOLOGIES

HUMAN BREAST TISSUE LYSATE

| | | | |
|------------------------|---|--|--------|
| Catalog Number: | Extraction 1, soluble protein fraction | | |
| | T2-011-T-1 | Human breast tumor tissue lysate | 100 µg |
| | T2-011-N-1 | Human breast normal tissue lysate (matched) | 100 µg |

| | | | |
|--|---|--|--------|
| | Extraction 2, insoluble protein fraction | | |
| | T2-011-T-2 | Human breast tumor tissue lysate | 100 µg |
| | T2-011-N-2 | Human breast normal tissue lysate (matched) | 100 µg |

Diagnosis: Infiltrating Ductal Carcinoma, grade 2, stage IIB. T₃N₀M₀

Sex / Age: Female, age 46.

Concentration: 1 mg/ml, 100 µg/vial.

The vial is provided with a 10% overflow. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides.

Storage: Aliquot single use volumes to avoid repeated freeze/thaw cycles. From time of receipt, this product is stable for 3 months at -20°C, or 12 months at -70°C.

Lysate Preparation: Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

| | | | |
|------------------------------|--------------------------------------|---------------------|-----------|
| Extraction 1: | PBS, pH 7.4 | 1 µg/ml Aprotinin | 1 mM NaF |
| Modified RIPA Buffer: | 1 mM EDTA | 1 µg/ml Pepstatin-A | 0.1% SDS |
| | 0.25% Na deoxycholate | 1 µg/ml Leupeptin | 1 mM PMSF |
| | 1 mM Na ₃ VO ₄ | | |

Extraction 2: PBS, pH 7.4, 5.0 M Urea, 2.0 M Thiourea, 50mM DTT, 0.1% SDS

Application: These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required.

Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays.

This material has tested negative for HbsAg, HIV 1/2, and HCV. Use *UNIVERSAL PRECAUTIONS* when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately.

Source: Integrated Laboratory Services-Biotech (ILSbio), Chestertown, MD 21620 www.ilsbio.com
ILS-7138

For Research Use Only



PROTEIN
BIOTECHNOLOGIES

PATHOLOGY REPORT

Catalog No. T2-011

Tissue: Breast

Location: Right breast.

Diagnosis: Infiltrating ductal carcinoma.

Stage: IIB T₃N₀M₀

Grade: II

Sex: Female

Age: 46 years

Gross findings: Tumor size 5 cm., ill demarcated.
Cut section firm and white.
Ipsilateral axillary lymph nodes examined: 4

Microscopic findings: The tumor is composed of proliferation of malignant epithelial cells. The epithelial cells are arranged as groups which have invaded the stroma. The tumor cells have irregular, basophilic nuclei and abundant cytoplasm. Nuclear chromatin is coarse with prominent nuclei. Mitotic figures are evident. The stroma is infiltrated by small numbers of chronic inflammatory cells. Necroses are seen but blood vessel invasions are not revealed. The lymph nodes show negative (0/4).